

WATER MANAGEMENT FOR SUGARBEET AND DRY BEAN

C. Dean Yonts
Extension Irrigation Specialist
Panhandle Research and Extension Center
University of Nebraska
Scottsbluff, NE 69361
Voice: 308-632-1246; Email: cyonts1@unl.edu

The past several years of sustained drought and expectations for below average snowpack and summer rains have many in agriculture searching for ways to stretch limited supplies of water. Not only has stream flow decreased, but ground water levels have declined and in many areas pumping restrictions have been imposed. At the same time, competition for water outside of agriculture further increases the demand for limited resources. The combination of drought and the increased demand for water will impose even more challenges for irrigated agriculture. It will require changing current irrigation practices and incorporation of new ideas to better utilize available water supplies as efficiently as possible. This means not only using irrigation water efficiently, but also using precipitation and stored soil water for crop production. Understanding the water needs of a crop will be a key to effective water management.

Water Use

The amount of water needed for irrigation varies by the crop being grown and the climatic conditions from year to year. Given in Table 1 are estimated water use rates for regionally grown crops.

| Alfalfa | Corn | Drybean | Spring Grain | Soybean | Sunflower | Sugarbeet | Winter Grain |
|---------|-------|---------|--------------|---------|-----------|-----------|--------------|
| 31-33 | 23-26 | 15-16 | 18-20 | 18-20 | 18-26 | 23-25 | 18-22 |

Table 1. Seasonal crop water use (in.) for regionally grown crops.

The depth from which sugarbeets get most of their water is generally considered to be from the top 3 to 4 ft of the soil profile. Sugarbeets use approximately 24 inches of water during the growing season and are often considered a crop that uses a large amount of water. Yet as we look closer, some of the crops we thought used less water, for example sunflowers and winter wheat, we find can use as much water as sugarbeets. However in the case of sunflowers and winter wheat, these crops can extract more water from the profile than most crops without adversely impacting yield potential. Sunflowers also have the ability to effectively extract water to depths of up to eight feet. In this case sunflowers may be viewed as a “drought tolerant” crop when in fact the crop has actually extracted more water from the soil and extracted water from deeper in the soil

profile. Anyone growing sunflowers knows that following this crop the soil can be left in a very dry condition the following spring.

Dry beans use approximately 16 inches of water during the growing season, which is approximately 8 inches less than what corn needs. This makes dry beans a good crop to grow if irrigation water is limited or if used as part of a crop rotation system to reduce overall irrigation needs. Dry beans are a shallow rooted crop with the majority of roots found in the top 18 in. of the soil profile. Roots can grow deeper into the soil profile to get water but this usually occurs late in the growing season as the plants begin to mature.

Water Management

The question of when is the best time to apply water to a crop often comes up when water supplies are limited. Some producers feel that stressing dry beans early in the growing season has little impact on yield and may even improve yield by forcing the roots to grow deeper into the soil profile. A similar question asked at the end of the season is whether stopping irrigation late in the season reduces yield?

For dry beans, early and late season water stress experiments have been conducted at the Panhandle Research and Extension Center in Scottsbluff, NE. The results of those experiments are given below.




Figure 1a. Effect of early season water stress on dry bean yield using sprinkler irrigation.




Figure 1b. Effect of early season water stress on dry bean yield using furrow irrigation.




Figure 2a. Effect of late season water stress on dry bean yield using sprinkler irrigation.




Figure 2b. Effect of late season water stress on dry bean yield using furrow irrigation.

Figures 1a and 1b, show the results of dry bean yield when water is limited during early season growth for sprinkler and furrow irrigation systems, respectively. The no stress treatment had irrigation starting approximately the last week in June to the first week in July. For the limited and high stress treatments, the initial irrigation was delayed for one week and two weeks, respectively. When sprinkler irrigation was used, yield tended to decline more as water stress increased compared to the furrow irrigation system. This is especially true for the high stress treatment under sprinkler. Yield loss was greater when water was withheld for two weeks because of the inability of the sprinkler system to replace soil water and meet the future water demand of the crop. A furrow irrigation system tends to refill the soil profile and is thus able to provide adequate water for future water use.

In figures 2a and 2b, the results of shutting off water late in the season are also shown for both sprinkler and furrow irrigation systems. The no stress treatment had irrigations throughout the growing season. Starting August 10, the limited stress treatment received every other irrigation that was scheduled for the no stress treatment while the high stress treatment received no further irrigations. Similar to the early season water stress results, dry beans irrigated with a sprinkler system showed a slightly steeper decline in yield as water stressed increased. The decline in yield is again likely related to the inability of the sprinkler irrigation system to supply water in excess to the requirements of the crop. Once irrigation was reduced or stopped less water was available in the soil profile to meet crop demands.

When comparing the early and late season experiments, there is a steeper decline in dry bean yield when water stress occurs at the beginning of the season as compared to water stress late in the season. These results are probably not uncommon and could be expected for most crops. Early in the season plant root development is limited and therefore water stress can occur rapidly. The lack of water during initial stages of plant growth likely impacts the majority of the root system. Late in the growing season, roots are more developed and reach further into the soil profile. Therefore water stress late in the season will first impact roots high in the soil profile while those deep in the profile may continue to extract some water to meet the needs of the crop. Finally, because the plant is nearing maturity, the need for water is declining on a daily basis and the root system can more easily keep up with the needs of the plant as water in the profile slowly moves to replace the water used by the crop.

For sugarbeets, the most critical time period when irrigation can affect final yield is during germination and early plant development. Inadequate soil water for germination and emergence results in reduced plant populations which in turn reduce final yield. Water stress after plants have emerged can result in seedling desiccation. At the early growth stages when root development is minimal, water stress can result in plant death with only a few days of warm dry winds. Often times if soil water is not adequate and stress begins, it is difficult to replenish the

soil water in a timely fashion. Even with center pivot irrigation, adequate water must be applied otherwise a light application merely meets the days evaporation demand. It is important to have an adequate supply of water in the soil below the seedling which allows soil water to migrate upwards and meet demands of the young seedling. As the season progresses, adequate water should be available to allow the sugarbeet to develop a good root system for extracting water from the soil.

The impact of late season water stress on sugarbeets was also studied at the Panhandle Research and Extension Center for both sprinkler and furrow irrigation systems. In these experiments, irrigation was either limited or stopped starting in mid-August. The results are given in figures 3a and 3b and show a yield decline as water stress increased for sprinkler and furrow systems, respectively. The decline however, was not as great as what might be expected. If the sugarbeet is allowed to develop a extensive roots system and water is available in the soil profile, it is capable of retrieving water from depths greater than 3 to 4 ft. In a current experiment irrigation water is being withheld from sugarbeets from July 15 to August 15. Preliminary results indicate very little difference in yield between full irrigation and no irrigation during the treatment period. The results indicate that like wheat and sunflowers, sugarbeets can effectively extract water from depths much greater than 3.0 ft and perhaps sustain periods of water stress without adversely impacting yield.




Figure 3a. Effect of late season water stress on sugarbeet yield using sprinkler irrigation.




Figure 3. Effect of late season water stress on sugarbeet yield using furrow irrigation.

Based on the results of the dry bean and sugarbeet experiments, if water is limited and the irrigator has the ability to choose when water supplies can be used, the choice should be to use water early in the season. Reducing irrigation late in the season has a smaller impact on yield than reducing irrigation early in the season and risking more of a yield reduction.