

WATER SAVINGS FROM CROP RESIDUE MANAGEMENT

Norman L. Klocke, Water Resources Engineer
Randall S. Currie, Weed Scientist
Troy J. Dumler, Extension Economist
Kansas State University, SWREC, Garden City, KS
Email:nklocke@ksu.edu

INTRODUCTION

Corn growers who irrigate in the Great Plains face restrictions in water, either from lower well capacities or from water allocations, and/or rising energy costs. They need water management practices to maximize grain production. When there is not enough water available to produce full yields, the goal for water management is to maximize transpiration and minimize non-essential water losses. One avenue for reducing non-essential water use is to minimize soil water evaporation.

Evapotranspiration is the combination of a two processes, transpiration and soil water evaporation. Transpiration, water consumed by the crop, is essential for the plants and correlates directly with grain production. Non-productive soil water evaporation has little utility. Soil water evaporation rates from bare soil are controlled by two factors. When the soil surface is wet, atmospheric energy that reaches the ground drives evaporation rates (energy limited phase). As the surface dries, evaporation rates are limited by the movement of water in the soil to the surface. In sprinkler irrigation during the growing season, most of the evaporation results from the energy limited process because of frequent soil wetting. Crop residues insulate the surface from energy limited evaporation.

Crop residues which are left in the field have value for soil and water conservation during the following non-growing season and the growing season of the next crop. Crop residues that are removed from the field after harvest are gaining value for livestock rations, livestock bedding, and as a source of cellulous for ethanol production. The water conservation value of crop residues needs to be quantified so that crop producers can evaluate whether or not to sell the residues or keep them on their fields. Reducing soil water evaporation in sprinkler management is one of the values of crop residues. This project was designed to measure soil water evaporation with and without a growing corn crop.

For presentation at the Central Plains Irrigation Conference, Colby, KS, February 24-25, 2009. Reprinted from 2008 CPIA meeting, Greeley, CO.

OBJECTIVES

1. Determine the water savings value of crop residues in irrigated corn.
2. Measure soil water evaporation beneath crop canopy of fully and limited irrigated corn.
 - a. From bare soil.
 - b. From soil covered with no-till corn residue.
 - c. From soil covered with standing wheat residue.
3. Calculate the contribution of evaporation to evapotranspiration.
4. Quantify soil water evaporation from partially covered soil with no crop canopy.
5. Predict potential economic savings from reducing evaporation with residues.

METHODS

Soil water evaporation was measured beneath a growing corn crop during the summers of 2004, 2005, and 2006 at Kansas State University's Research and Extension Center near Garden City, Kansas. The soil at the research site was a Ulysses silt loam. Mini-lysimeters were used for the primary evaporation measurement tool. They contained undisturbed soil cores 12 inches in diameter and 5.5 inches deep. The soil cores were extracted by pressing PVC tubing into the soil with a custom designed steel bit. The PVC tubing became the sidewalls for the mini-lysimeters. The bottom of the cores were sealed with galvanized discs and caulking. Therefore, water could only escape from the soil by surface evaporation, which could be derived from daily weight changes of the mini-lysimeters. Weighing precision produced evaporation measurements with a resolution of ± 0.002 in/day.

Volumetric soil water content was measured bi-weekly in the field plots to a depth of 8 ft in 1 ft increments with neutron attenuation techniques. The change in soil water, from the start to the end of the sampling period, plus measurements of rainfall and net irrigation were the components of a water balance to estimate crop evapotranspiration (ET_c).

Measurements of crop residue coverage on the soil surface were adapted from line transect techniques. A coarse screen was laid over a mini-lysimeter. Observations of the presence or absence of residue were recorded for each intersection of screen material. The fraction of the presence of residue and total observations was converted into a percentage of coverage.

Four replications of bare, corn stover, or wheat stubble surface treatments were placed in high and low frequency irrigation treatments. High frequency irrigation was managed to meet atmospheric demand for full crop evapotranspiration (ET_c). The low frequency irrigation treatment received approximately half this amount in half the irrigation events.

An additional experiment was conducted to find the soil water evaporation rates from soil surfaces that were partially covered with crop residues. A controlled area was established for the experiment where the mini-lysimeters were buried in PVC sleeves at ground level, arranged adjacent to one another in a geometric pattern. Movable shelters were available to cover the mini-lysimeters during rain events but were open during other times. There was no crop canopy over the mini-lysimeters, which were surrounded by mowed, irrigated grass. The mini-lysimeters were weighed daily. Two irrigation treatments, that approximated the companion field study, were watered with 1 or 2 per hand irrigations per week. Partial surface cover treatments had 25%, 50%, and 75% of the surface covered with corn stover which was placed on the mini-lysimeters. Mini-lysimeters with 100% coverage from corn stover and 85% coverage with standing wheat stubble were the same configuration as the field experiment. Evaporation results were normalized with reference ET (ET_r) which was calculated with on-site weather factors and an alfalfa referenced ET_r model (Kincaid and Heermann, 1984).

RESULTS

Within Canopy Field Results

Soil surface cover on the mini-lysimeters was measured at the start of the growing season. Corn stover and standing wheat stubble completely covered the mini-lysimeters in 2004 (table 1). Corn stover continued to completely cover the mini-lysimeters in 2005 and 2006, but the wheat stubble coverage was 91-92% in those years. The 2004 and 2005 wheat crops were shorter in stature due to less fall growth. This led to less wheat stubble coverage of the mini-lysimeters during the following year.

Table 1. Crop residue percentage cover at the end of the growing season for mini-lysimeters in corn field plots during 2004-2006 near Garden City, Kansas.

Crop Residue Cover	Dry Matter tons/ac	Residue Coverage*
-----2004-----		
Bare	0.0	0
Corn	7.3	97
Wheat	9.8	98
-----2005-----		
Bare	0.0	0
Corn	9.5	100
Wheat	6.3	91
-----2006-----		
Bare	0	0
Corn	7.5	100
Wheat	4.3	92

*Percentage of soil surface covered by residue, determined by the modified line transect method.

When data from all years and water frequency treatments were combined, the effects of surface treatments could be isolated. Average soil water evaporation (Avg E) from the bare surface treatment was significantly more than Avg E from the two residue covered treatments (table 2). Wheat stubble surface coverage was than corn stover coverage in 2005 and 2006, resulting in more E with wheat stubble. Daily average ET_c and ET_r data were the same over all mini-lysimeters since the annual data was averaged over all irrigation treatments. Bare soil E for the Ulysses silt loam was 30% of ET_c, which was the same result as a study with Valentine fine sandy soils in west-central Nebraska (Klocke et al., 1985). E as a ratio of ET_c or ET_r showed that crop residues reduced E by 50% compared with bare soil. A similar study with silt loam soils in west-central Nebraska showed that bare soil E under a corn canopy during the growing season could be reduced from 0.07 inches/day to 0.03 inches/day by adding a mulch of wheat stubble lying flat on the surface with 100% surface coverage (Todd et al., 1991).

Differences in E between bare soil and residue treatments, which were 0.02-0.03 inch per day, may seem small; however, if these daily differences were extrapolated over a 110 day growing season, total differences in E would be 2.2-3.3 inches. Similarly, E as a fraction of ET_c was 0.30 for bare soil and 0.15-0.16 for the residue cover treatments. Growing season ET_c values for corn can be 24-26 inches in western Kansas. Using the values of E as a fraction of ET_c (table 2), potential water savings could be 3.7-4.0 inches with full soil surface coverage.

Table 2. Average soil water evaporation and evaporation as a ratio of crop evapotranspiration (ET_c) and reference ET (ET_r) for all bare soil and crop residue covered treatments under a corn crop canopy during 2004-2006 in Garden City, KS.

Surface Cover	Avg E in/day	ET _c in/day	E/ET _c *	ET _r in/day	E/ET _r
Bare	0.06a	0.23	0.30a	0.27	0.22a
Corn Stover	0.03c	0.23	0.15c	0.27	0.11c
Wheat Straw	0.04b	0.23	0.16b	0.27	0.12b
LSD _{.05} **	0.003		0.02		0.05

Means with same letters in the same columns are not significantly different for alpha=.05.

The influence of crop canopy shading canopy on soil water evaporation rates was observed by averaging data over years, surface cover treatments, and irrigation frequency treatments (table 3). Evaporation decreased as crop canopy and ground shading increased. The trend reversed as the crop matured and shading decreased. Concurrently, crop ET and reference ET increased from planting through mid-season and then decreased through the rest of the growing season. The ratio of Avg E to ET_c and ET_r declined during the growing season when the two factors were combined.

Table 3. Soil water evaporation (Avg E) and evaporation as a ratio of crop ET (ETc) and reference ET (ETr) during the growth stages of corn for all mini-lysimeter treatments during the 2004-2006 growing seasons at Garden City, KS.

Growth Stage	Avg Days In Growth Stage	Avg E	ETc	E/ETc	ETr	E/ETr
		in/day	in/day		in/day	in/day
Vegetative	28	0.06a	0.22b	0.27a	0.35	0.17a
Pollination	18	0.05b	0.27a	0.20b	0.33	0.15b
Seed Fill	30	0.03c	0.20c	0.15c	0.25	0.12c
LSD _{.05}		0.002	0.02	0.02		0.05

Means with same letters in the same columns for the same year are not significantly different for alpha = 0.05.

Partial Cover Results from Control Area

Even though average daily evaporation rates among the bare and 25%, 50%, and 75% residue covered treatments could be measured and were significantly different from one another, the magnitudes of these differences were small (table 4a). The 100% covered treatment with corn stover and the standing wheat stubble with 85% cover produced significantly less E than the other treatments. Lateral heat flow from the bare portion of the partially covered surface could have caused increased surface temperatures under the corn stover. Similarly, soil water could move from under partially covered surface to the bare portion of the surface, increasing E (Chung and Horton, 1987).

Crop residues that were distributed across the surface, needed to cover more than 75-80% to have an effect in reducing E when there was no crop canopy. Nearly complete surface coverage influenced E nearly the same with and without crop canopy.

Table 4. Soil water evaporation during Spring and Fall 2005 and Fall 2006 for full and partial crop residue surface covers at Garden City, Kansas.

a. Surface Cover	Avg E	E/ETr*
	--in/day--	
Bare 0%	0.08a	0.26a
Corn 25%**	0.07b	0.25b
Corn 50%	0.07c	0.24c
Corn 75%	0.07a	0.26a
Corn 100%	0.04e	0.14e
Wheat 85%	0.05d	0.18d
LSD _{.05}	0.002	0.005

Means with same letters in the same columns for the same variable are not significantly different at alpha = 0.05.

ECONOMIC IMPACT

Crop residues can also have an effect on non-growing season. A field study in eastern Colorado during October-April of the years 2000-2004 showed that corn residues increased stored soil water by 2 inches when compared with conventional stubble mulch tillage in dryland management (Neilson, 2006). Dryland studies in Nebraska have demonstrated that wheat stubble increased non-growing season soil water storage by 2-2.5 inches when compared with bare soil (Klein, 2007).

The water savings from crop residues can have one of three impacts on income. First, if irrigation is applied in excess of water requirements of the crop in a no-till system, there could be no economic benefits from the crop residues. The excess water could leach past the root zone with no value to crop production. Second, if water supplies are adequate to grow a fully irrigated crop, pumping costs can be reduced by the difference between tilled and no-till management. Irrigators in this situation need to monitor soil water during the growing season to find the reduction in irrigation needed from crop residue management and time irrigations accordingly. Third, if the irrigation system cannot keep up with crop water requirements, the crop may be under water stress all or part of the growing season. Water savings from crop residues in no-till management can be transferred from bare soil evaporation losses to water that can be used by the crop (transpiration) for better yield returns. In this case there would be no change in irrigation pumping.

Irrigation requirements and production costs vary from year-to-year and from one irrigator to another. Commodity prices also vary from year-to-year. As demonstrated in this study, nearly full coverage of the soil surface was needed to reduce soil water evaporation and reap benefits from the crop residues. The following is one example of economic impacts on income for irrigated corn where growing season and non-growing season crop residue management combines for saving 3-5 inches of water annually:

Situation 1. Irrigation applications in excess of crop needs can lead to soil water leaching below the root zone and there are no benefits from the crop residues.

Situation 2. Irrigation requirements are reduced for a fully irrigated crop from crop residue management where pumping is reduced to account for less irrigation needs.

Pumping costs = \$9 per acre for each inch pumped
Total savings for 3-5 inches less water pumped = \$27-45 per acre

Situation 3. The irrigation system cannot provide enough water to meet the full water requirements of the crop. Three to five inches of water savings from crop residue management could shift soil water evaporation to transpiration. Corn

yields increase 12 bushels per acre for each inch of water that is transferred from evaporation to transpiration. When corn price is \$4.50 per bushel, 3-5 inches of water savings from reduced evaporation would produce \$162-\$270 per acre additional income.

Additional growing and non-growing season benefits from crop residues include capturing precipitation, enhancing infiltration, reducing runoff, and reducing soil erosion. All of these benefits have economic value for crop production and land values, but they are more difficult to measure than direct water conservation effects of crop residue management.

ACKNOWLEDGEMENTS

Partial funding for this research was furnished by: the USDA-ARS Ogallala Aquifer Project and the US Department of Interior.

REFERENCES

Chung, S., and R. Horton. 1987. Soil heat and water flow with a partial surface mulch. *Water Res. Res.* 23(12):2175-2186.

Kincaid, D.C., and D.F. Heermann. 1984. Scheduling irrigation using a programmable calculator. NC-12. Washington, D.C.:USDA-ARS.

Klocke, N.L., D.F. Heermann, and H.R. Duke. 1985. Measurement of evaporation and transpiration with lysimeters. *Trans. of the ASAE.* 28:1:183-189 & 192.

Klein, R.N. 2008. Improving you success in no-till. In Cover Your Acres Proceedings. Kansas State Research and Extension. Jan. 23-23, 2008. Oberlin, KS. p. 22-26.

Neilsen, D.C. 2006. Crop residue and soil water. In Central Plains Irrigation Conference & Exposition Proceedings. Feb. 21-22, 2006. Colby, KS. p. 136-139.

Todd, R.W., N.L. Klocke, G.W. Hergert and A.M. Parkhurst. 1991. Evaporation from soil influenced by crop shading, crop residue and wetting regime. *Trans. of the ASAE.* 34:2:461-466.